

Comment on "Spectral Signatures of the Fulde-Ferrell-Larkin-Ovchinnikov Order Parameter in One-Dimensional Optical lattices"

In a recent letter Reza Bakhtiari *et al.* [1] studied an imbalanced two-component atomic Fermi gas in a one-dimensional (1D) optical lattice with a trapping potential, within the Bogoliubov-de Gennes (BdG) approximation. They showed that the prominent oscillations of the pairing gap (within the BdG approximation), characteristic of a Fulde-Ferrell-Larkin-Ovchinnikov state (FFLO), could be detected in the rf spectra and in the momentum-resolved photoemission spectra of the gas. In this comment we show that the BdG approximation not only produces inaccurate results for the examples presented in [1], but that they are qualitatively incorrect making the analysis of the rf spectra unreliable and shedding doubts on the applicability of rf spectroscopy to detect the FFLO state in 1D optical lattices.

The Letter [1] is devoted to 1D optical lattices without any reference to higher dimensions where the BdG approximation could be valid. The use of the BdG approximation is justified by saying that it provides qualitative information on the system and allows one to calculate the rf spectrum. In spite of the known failure of the BdG approximation in 1D systems, it is still used [2]. In the absence of a trap the BCS treatment gives large errors in the order parameter as compared to the exact solution (see Ref. 19 of the letter). With the inclusion of the trap we have to resort to the Density Matrix Renormalization Group (DMRG) to produce numerically "exact" solutions for 1D lattice problems of moderate sizes (~ 100 sites). Indeed, the DMRG has already been used to study the FFLO phase in a 1D trapped lattice gas (see Ref. 11 and 16 of the letter). Therefore the DMRG constitutes an ideal benchmark to test the accuracy of the BdG approximation in 1D lattice problems.

In Fig. 1 we show the densities and the absolute value of the gap for a system with polarization $P = 0.23$ ($N_\uparrow = 40$, $N_\downarrow = 25$) in the upper panels and for $P = 0.70$ ($N_\uparrow = 40$, $N_\downarrow = 7$) in the lower panels, in a lattice of $L = 150$ sites. Panels A and D correspond to the BdG approximation neglecting the Hartree term like in Fig. 1 of [1]. In panels B and E include the Hartree term into the BdG approximation. Panels C and F are DMRG results. The gap parameter is defined as $|\Delta_i| = U\sqrt{\langle n_{i\uparrow}n_{i\downarrow} \rangle - \langle n_{i\uparrow} \rangle \langle n_{i\downarrow} \rangle}$, where the expectation values are taken in the ground state wavefunctions of the three approximations. Within the BdG approximations this definition coincides with local pairing gap. The Hartree term, neglected in [1], does modify the results towards the "exact" DMRG results. However, both BdG approximations are still quite far from the DMRG results, specially for lower polarizations. The most dramatic differences can be seen in the pairing gap with very large amplitude oscillations which are completely softened in the DMRG results. It might be argued




FIG. 1: (Color online) Density of majority and minority atoms n_i and absolute value of the local gap parameter $|\Delta_i|$ as a function of the site index i . Upper panels show results for $P = 0.23$ and lower panels for $P = 0.70$. From left to right we show results of BdG without the Hartree term, BdG with the Hartree term and DMRG.

that the dimensions of the systems considered are small, however, doubling the size of the systems the authors found similar results as we did using DMRG.

The subsequent analysis of the rf spectra is based on the BdG results of panels A and D. Taking into account that the gap oscillation are reduced by one order of magnitude in the DMRG results, it is doubtful that rf spectroscopy could provide information about the spatial structure of the pairing gap. Whether rf spectroscopy could signal the FFLO phase in 1D lattice systems is still an open question which might be confirmed by rf experiments, or addressed numerically by means of a DMRG study. Note that within the Correction Vector Approach the calculation of the excitation spectrum is not needed.

This work is supported by Spanish grants FIS2006-12783-C03-01 and CAM-CSIC CCG07-CSIC/ESP-1962.

R. A. Molina¹, J. Dukelsky¹, and P. Schmitteckert²
¹Instituto de Estructura de la Materia, CSIC, Serrano 123, 28006 Madrid, Spain
²Institut für Nanotechnologie, Forschungszentrum Karlsruhe, 76021 Karlsruhe, Germany

PACS numbers: 03.75.Ss, 74.45.+c, 78.90.+t

-
- [1] M. Reza Bakhtiari, M. J. Leskinen, and P. Törmä, Phys. Rev. Lett. **101**, 120404 (2008).
[2] T. K. Koponen, T. Paananen, J. P. Martikainen, M. R. Bakhtiari, and P. Törmä, New J. Phys. **10**, 045014 (2008); X. Liu, H. Hu, and P. D. Drummond, Phys. Rev. A **78**, 023601(2008) (Ref. [11] of the letter).